

# **Guidelines for a Banded Pesticide Tax Scheme, Differentiated According to Human Health and Environmental Risks**

Compiled by  
Erlend Spikkerud, Terje Haraldsen, Abdelkarim Abdellaue and Marianne T. Holmen  
Norwegian Food Safety Authority, National Centre of Plants and Vegetable Foods

January 2005

## Contents

<b>1. BACKGROUND.....</b>	<b>3</b>
<b>2. NEW TAX BANDS.....</b>	<b>3</b>
<b>3. TAX BURDEN.....</b>	<b>4</b>
<b>4. CALCULATING THE TAX.....</b>	<b>5</b>
<b>5. HUMAN HEALTH .....</b>	<b>5</b>
5.1. INTRINSIC PROPERTIES.....	5
5.2. EXPOSURE .....	6
5.3. TOTAL SCORE FOR HUMAN HEALTH.....	7
5.4. CATEGORIES FOR HUMAN HEALTH .....	7
<b>6. ENVIRONMENT.....</b>	<b>8</b>
6.1. TOTAL SCORE FOR ENVIRONMENT .....	8
6.2. SCORE FOR THE INDIVIDUAL ORGANISMS/PARAMETERS .....	9
6.3. CHOICE OF PARAMETERS .....	12
6.4. CATEGORIES FOR ENVIRONMENT.....	12

## 1. Background

As a follow-up of the Action Plan for Reduced Risk of Pesticide Use (1998-2002), a banded tax scheme for pesticides was introduced in 1999, based on differentiated human health and environmental risks. An internal committee in the Norwegian Agricultural Inspection Service assessed this system in 2003. In a report (Landbrukstilsynet 2003a), the committee concluded by making the following suggestions for improvements:

- Adjust the criteria for the tax bands with regard to health effects, in accordance with the health risk indicator.
- Include the exposure to pesticides during the handling of treated plant material in the part of the tax scheme related to human health.
- Base the tax bands for seed dressings on human health hazard, since exposure and the substances' intrinsic properties can vary significantly.
- Include relevant metabolites in the health classification. In some cases, metabolites may present a greater health risk than the active ingredients.
- Change the guidelines for environmental aspects of the banded tax scheme, to a system that takes more consideration to the total environmental hazard of pesticide use.
- Revise the guidelines for determining the standard area dose (maximum application rate in the or those crop(s) accounting for 50 % or more of the product's total application), in order to improve their user-friendliness.
- Increase the number of health and environmental categories for products used by certified applicators. This will lead to an improved differentiation between products with low or high risk, and less differences between categories. This should perhaps result in a larger number of tax bands.

This report takes the above-mentioned items into consideration. For health-related issues, the report is only a revision and an adjustment of the previous guidelines (Landbrukstilsynet 2003b). For environmental issues, a whole new system was devised, which resembles a revised version of the risk indicator scheme. The revised guidelines for standard area doses are published in a separate report (Planteforsk 2003).

## 2. New tax bands

Each product is assigned to one of three health categories and one of three environmental categories. Previously, there were only two categories for each. Based on this classification, products are assigned to one of the tax bands presented in the table below. These tax bands form the basis for the calculation of environmental taxes.

**Tax band 1 (0.5 x base rate):**

Products with a low health and environmental risk.

**Tax band 2 (3 x base rate):**

Products with a low health risk and medium environmental risk, or medium health risk and low environmental risk.

**Tax band 3 (5 x base rate):**

Products with a low health risk and high environmental risk, or medium health risk and medium environmental risk, or high health risk and low environmental risk.

**Tax band 4 (7 x base rate):**

Products with a high health risk and medium environmental risk, or medium health risk and high environmental risk.

**Tax band 5 (9 x base rate):**

Products with a high health and environmental risk.

**Tax band 6 (50 x base rate):**

Concentrated consumer products.

**Tax band 7 (150 x base rate):**

Ready-for-use consumer products.

The total number of tax bands remains unchanged at seven, but the number of bands for products used by certified applicators is increased from three to five in order to enable a better differentiation. The number of bands for consumer products (previously numbers 5 and 6, now 6 and 7) remains the same. The former tax bands 0 for adjuvants and 1 for seed dressings are removed. Adjuvants will remain to be exempt from pesticide taxes, and are not included in the scheme. Seed dressings and microbiological agents are placed in the lowest environmental category, but can be moved to a higher health category depending on the product's health risk.

### 3. Tax Burden

A condition for the reorganisation of the environmental tax scheme was that the total tax burden should not be changed. The factors for the various tax bands are thus set to achieve this objective. The base tax rate remains the same at NOK 25 per hectare. For products used by certified applicators, the tax for products with a low health and environmental risk is halved (factor reduced from 1 to 0.5), while the tax for products with a high health and environmental risk is raised by 12.5 % (factor increased from 8 to 9.). The tax for consumer products has not been changed.

## 4. Calculating the Tax

The environmental tax is acreage-based. The tax for a specific pesticide per hectare (ha) is calculated by multiplying the base rate (NOK 25) with the factor for the tax band in question (0.5 - 150):

$$\text{Tax per hectare} = 25 \times \text{factor for tax band}$$

This is converted to tax per kg or litre by multiplying with 1000 and dividing by the standard area dose (grams or ml/ha):

$$\text{Tax per kg or litre} = 25 \times \text{factor} \times 1000 / \text{standard area dose}$$

## 5. Human Health

When assigning products to the three health categories, consideration is taken to intrinsic properties, exposure during mixing and exposure during spraying.

### 5.1. Intrinsic properties

Based on the products' intrinsic properties, they are classified into four groups for potential health hazard, in accordance with the health hazard labelling. The following categories of danger are used: Very toxic (T+), toxic (T), harmful (Xn), corrosive (C) and irritant (Xi).

#### Group 1: Products with a low potential health hazard:

- Products without risk-phrases (R-phrases) for health hazard.

#### Group 2: Products with a medium potential health hazard:

- Harmful (Xn) products not included in group 3.
- Products classified as irritant (incl. allergy) (Xi).
- Products containing allergenic substances which must be declared on the product label.
- Products labelled:
  - R33 Danger of cumulative effects
  - R64 May cause harm to breastfed babies

#### Group 3: Products with a high potential health hazard:

- Products labelled very toxic (T+) or toxic (T)
- Products labelled corrosive (C)
- Products that are harmful (Xn), and are labelled with one of the following:
  - R68 Possible risk of irreversible effects
  - R48 Danger of serious damage to health by prolonged exposure
  - R62 Risk of impaired fertility

- R63 Possible risk of harm to the unborn child
- R40 Limited evidence of a carcinogenic effect

**Group 4: Products with a very high health hazard:**

- Products marked with two or more of the following R-phrases:
  - R23 Toxic by inhalation
  - R24 Toxic in contact with skin
  - R25 Toxic if swallowed
  - R26 Very toxic by inhalation
  - R27 Very toxic in contact with skin
  - R28 Very toxic if swallowed
  - R39 Danger of very serious irreversible effects
  - R40 Limited evidence of a carcinogenic effect
  - R45 May cause cancer
  - R46 May cause heritable genetic damage
  - R48 Danger of serious damage to health by prolonged exposure
  - R49 May cause cancer by inhalation
  - R60 May impair fertility
  - R61 May cause harm to the unborn child
  - R62 Risk of impaired fertility
  - R63 Possible risk of harm to the unborn child
  - R68 Possible risk of irreversible effects

## **5.2. Exposure**

Users are typically exposed to pesticides during such operations as mixing and diluting, application (often spraying) and handling treated crops. Exposure during mixing depends on the concentration of the active ingredient in the product, the number of dilutions and the type of formulation. The latter is of major importance. The degree of exposure during application depends on such factors as the concentration of active ingredient in the diluted liquid, treatment dose, treated area and especially the application method.

The tax scheme takes the type of formulation and application method into consideration. Mixing is given more weight, since users are working with concentrated products when mixing, and pesticides are easily spilled.

With regard to exposure during mixing, products can be divided two groups depending on the type of formulation: low exposure (factor 1) and high exposure (factor 3).

- Examples of products that can give low exposure during mixing: Seed dressings used in approved facilities, ready-to-use solutions, granules, water-soluble pellets and water-soluble bags.
- Examples of products that can give high exposure during mixing: Emulsions, powders, suspensions and other soluble concentrates.

With regard to exposure during application, products can also be divided in two groups, depending on the application method: low exposure (factor 1) and high exposure (factor 2). The method of application depends on such factors as the type of crop in which the pesticide is used. For products approved for several crops, the same method as for the calculation of standard area doses is used, i.e., the or those crops accounting for 50 % or more of the product's total application. In cases where there are several major crops, the selected application method is based on the crop with the highest potential degree of exposure.

- Examples of application methods that can give low exposure: Seed treatment in approved facilities, mist blowers, fumigation, granular pesticide application, tunnel spraying in forest nurseries.
- Boom field or field crop sprayers are classified in relation to their operating pressure. High pressure gives small particles, which are easily stirred up, thus resulting in a high degree of pesticide exposure among applicators. Such assessments are made on a case-to-case basis.
- Examples of application methods that can give high exposure: Knapsack (mist) sprayers, tractor-mounted mist sprayers, manual pressure sprayers.

### 5.3. Total score for human health

Based on the pesticide's intrinsic properties and the considerations to exposure presented above, the total scores for human health are presented in the table below. The total exposure is obtained by multiplying the factors for "during mixing" and "during spraying". The total score for human health (H) is obtained by multiplying total exposure by the factor for intrinsic properties. The respective factors are shown in parentheses.

Intrinsic properties	Total exposure			
	<i>During mixing</i>			
	Low (1)		High (3)	
	<i>During spraying</i>		<i>During spraying</i>	
	Low (1)	High (2)	Low (1)	High (2)
Low (1)	1	2	3	6
Medium (2)	2	4	6	12
High (4)	4	8	12	24
Very high (8)	8	16	24	48

### 5.4. Categories for human health

Based on the system described above, products are classified into three health categories:

Health category	Total score for human health
Low	<8
Medium	8-16
High	>16

Exceptions:

- Greenhouse pesticides or seed dressings that are acutely toxic in contact with skin or by inhalation (T or T+), or that present a danger of serious irreversible effects, or present a risk of cancer, impaired fertility, harm to the unborn child or genetic damage (R39, R40, R45, R 46, R48, R49, R60, R61, R62, R63, R68). These products are to be classified in the 'medium' health category, due to the risk of damage to human health during handling of treated plant material or treated seed, and when cleaning seed treatment machinery.
- Products with active ingredients that are degraded to metabolites which may cause irreversible damage, cancer, impaired fertility, harm to the unborn child or genetic damage (R39, R40, R45, R 46, R48, R49, R60, R61, R62, R63, R68). (This applies to cases in which the hazard classification of metabolites is not shown in the labelling of the active ingredient). These products are to be classified in the 'medium' health category.

## 6. Environment

### 6.1. Total score for environment

The total score for environment is calculated for each active ingredient in each product in accordance with the equation below. In other words, a substance found in several different products will achieve different scores in the various products, depending on the respective dosage and application.

$$\text{Total score for environment} = \text{Te} + \text{Ta} + \text{Tb} + \text{A} + \text{L} + \text{P} + \text{B} + \text{F}$$

Te	= Score for earthworms
Ta	= Score for bees and other arthropods
Tb	= Score for birds
A	= Score for aquatic organisms
L	= Score for leaching potential
P	= Score for persistence
B	= Score for bioaccumulation
F	= Score for formulation type

For products containing several active ingredients, the highest score for each of the above-listed parameters is used to determine the product's total score.

In case of missing or insufficient documentation, a high score can be assigned to a substance.

Certain products have so specific areas of use or properties, that the equation is not suited for use. Examples hereof include seed dressings only used in approved facilities, greenhouse pesticides and microbiological products. The environmental hazard of these



products is considered to be so insignificant, that they are given a score of 0 without any further calculations.

## 6.2. Score for the individual organisms/parameters

### Earthworms (Te)

The European Union's working group FOCUS (1997) proposed a simple model for the estimation of pesticide concentration in the soil immediately after spraying (predicted initial environmental concentration, PIEC). 50 % plant cover is chosen as a standard scenario. The toxicity exposure ratio (TER) for earthworms is thereafter calculated by dividing toxicity (14-day LC<sub>50</sub>) by PIEC.

For earthworms, the European and Mediterranean Plant Protection Organisation (EPPO) defined a threshold value of 100 for acute TER, whereas the EU (Uniform Principles) defined a threshold value of 10. On the basis of the TER, the Te score (Terrestrial earthworm) is determined, according to the table below:

TER based on 14d LC <sub>50</sub>	Te
>100	0
10-100	2
<10	4

### Bees and other arthropods (Ta)

For bees, EU's Uniform Principles define Hazard Quotients for oral exposure (Qho) and for contact exposure (Qhc), as dose (g/ha) divided by oral or contact toxicity (LD<sub>50</sub>, µg/bee), respectively. The threshold value is set at 50. Based thereupon, the highest hazard quotient is used to determine the Ta-score (Terrestrial – arthropod), according to the table below. Two additional points are added if the product is classified “toxic for insects” in accordance with separate labelling criteria.

Qho or Qhc	Ta without labelling for insect toxicity	Ta with labelling for insect toxicity
<50	0	2
50-1000	1	3
>1000	2	4

### Birds (Tb)

In the EU's (2002) guidance document for birds and mammals, several steps are defined for risk assessment of birds. Here, we only use step 1, which is a standardised, realistic “worst case” with a standard exposure scenario for herbivorous birds. We selected medium-sized birds with a body weight of 300 g as the standard. The bird's daily intake of a substance is given by the following equation:

$$\text{ETE} = \text{FIR}/\text{body weight} \times \text{C}$$

ETE = Estimated daily intake

FIR = Food Intake Rate

C = Concentration of the substance in the diet (mg/kg)

According to Table 4 of the guidance document, the FIR/body weight ratio for the above-mentioned standard bird size is 0.76. RUD (residue unit dose) is used to calculate pesticide residues in the diet from dosage per unit area. In the table, RUD is set at 87 for our scenario. To obtain the concentration C (mg/kg), RUD is multiplied with dosage as kg/ha. Thus, we obtain the following equation:

$$\text{ETE} = 0.76 \times 87 \times \text{dose} = 66.12 \times \text{dose}$$

ETE corresponds to PIEC, which is used to calculate TER by dividing acute oral toxicity ( $\text{LD}_{50}$ ) by PIEC. For birds, EU's Uniform Principles define a threshold value for TER of 10 from acute and sub-acute studies. The lowest TER value, determined for the most sensitive species, is used to define the Tb score (Terrestrial bird), according to the table below:

TER	Tb
>10	0
5-10	1
1-5	2
0.1-1	3
<0.1	4

### **Aquatic organisms (A)**

Sources of pesticides in surface water are mainly spray drift, surface runoff or discharge via drainage systems. The EU working group FOCUS (2002) recommended three steps for the calculation of pesticide concentrations in surface water and sediments. We have chosen to use step 1, which combines spray drift, surface runoff, erosion and/or drainage, and use these parameters to estimate the predicted initial environmental concentration (PIEC). To prevent the scheme from becoming too complex, we chose to disregard repeated treatments, and only to consider the aquatic phase (not sediments).

In connection with pesticide registration, effect studies of algae, pelagic invertebrates (*Daphnia*) and fish must be submitted. In addition, studies on aquatic plants and sediment invertebrates may have been submitted. TER values for algae/aquatic plants and daphnia/fish are determined by dividing acute toxicity ( $\text{LC}_{50}/\text{EC}_{50}$ ) by PIEC from the FOCUS-scenario described above.

The EU's Uniform Principles define threshold values for TER. For acute studies of daphnia and fish, the threshold value is 100. For chronic studies and algae/aquatic plant

studies, the threshold is 10. On this basis, one score is determined for algae/aquatic plants and one for daphnia/fish in accordance with the table below. The highest of these scores is used as the score for aquatic environment (A).

TER based on studies on algae or aquatic plants	TER based on acute studies on daphnia or fish	A
>10	>100	0
1-10	10-100	1
0.1-1	1-10	2
0.01-0.1	0.1-1	3
<0.01	<0.1	4

### Leaching (L)

The SCI-GROW model (EPA 2001) calculates a score for leaching risk, based on dosage, mobility (Koc) and persistence (DT<sub>50</sub>). This is used to assign a score L for leaching in accordance with the table below:

SCI-GROW	L
<5	0
5-20	1
20-100	2
100-500	3
>500	4

Substances which are frequently detected in environmental monitoring programmes can be given a higher score than indicated by their inherent properties.

### Persistence (P)

Based on half-life in soil, each substance is assigned a persistence factor (P) in accordance with the table below.

DT <sub>50</sub> (days)	P
<10	0
10-45	1
45-90	2
90-180	3
>180	4

### Bioaccumulation (B)

Based on log P<sub>ow</sub> and half-life in soil, a score for bioaccumulation (B) is assigned in accordance with the table below.

Log P <sub>ow</sub>	Half-life in soil (days)			
	<10	10-90	90-180	>180
<3	0	0	0	0
3-4	0	0	4	2
>4	1	2	3	4

### Formulation type (F)

In order to take the spillage risk during the mixing of pesticides into consideration, a score for type of formulation (F) is assigned in accordance with the table below.

Formulation type	F
Water-soluble bags, water-soluble pellets, granules	0
Emulsions, powders, suspensions and other soluble concentrates	1

### 6.3. Choice of parameters

*Dose:* Standard area dose is used as dosage in all calculations.

*Crop:* The crop(s) on which the determination of the standard area dose is based.

*Toxicity:* The value for toxicity (LC<sub>50</sub>/LD<sub>50</sub>/EC<sub>50</sub>) is based on the most sensitive standard species in laboratory studies. When not available, results from other submitted studies may be used.

*Adsorption in soil:* The average of all results from standard laboratory studies is used to determine K<sub>oc</sub>. Results from soil types which clearly are not relevant for Norwegian conditions, e.g., with extremely high contents of organic matter, may be omitted.

*Persistence in soil:* The average of all results from standard laboratory studies at room temperature is used to determine DT<sub>50</sub>. Results from soil types which clearly are not relevant for Norwegian conditions, may be omitted.

### 6.4. Categories for environment

Based on the products' total score they are assigned to one of three environment categories in accordance with the table below.

<b>Environment category</b>	<b>Score for environment</b>
Low	<4
Medium	4-8
High	>8

Exceptions:

- Products that, based on a general assessment, clearly represent a low environmental risk can be directly placed in the lowest environment category.
- Products with incomplete documentation, but which are suspected to represent a high environmental risk, can be classified as "medium" or "high".
- Products can be placed in a higher category than indicated by their active ingredient(s), if metabolites represent a higher environmental risk than the active ingredient(s).

## Literature

EPA 2001. [http://www.epa.gov/oppefed1/models/water/scigrow\\_description.htm](http://www.epa.gov/oppefed1/models/water/scigrow_description.htm)

EU 2002. Guidance Document on Risk Assessment for Birds and Mammals Under Council Directive 91/414/EEC. Sanco/4145/2000 – final. 25. September 2002.

FOCUS 1997. Soil persistence models and EU registration. The final report of the work of the Soil Modelling Work group of FOCUS (Forum for the Co-ordination of pesticide fate models and their Use).

FOCUS 2002. FOCUS surface water scenarios in the evaluation process under 91/414/EEC. Report prepared by the FOCUS Working Group on Surface Water Scenarios. Final Draft 06-06-02.

Landbrukstilsynet 2000. Målemetoder for risiko ved bruk av pesticides. Versjon 3. Mai 2000.

Landbrukstilsynet 2003a. Evaluering av avgiftssystemet for pesticides.

Landbrukstilsynet 2003b. Retningslinjer for klassifisering av pesticides i avgiftsklasser differensiert etter helse- and miljøegenskaper. Versjon 1.2. Revidert mars 2003.

Planteforsk 2003. Retningslinjer for fastsettelse av normerte arealdoser (NAD) for pesticides. Versjon 2.0. Revidert 10/2003.